

Name of research institute or organization:

**Laboratory of Atmospheric Chemistry, Paul Scherrer Institute,
CH-5232 Villigen PSI, Switzerland**

Title of project:

The Global Atmosphere Watch Aerosol Program at the Jungfrauoch

Project leader and team:

Prof. Dr. Urs Baltensperger, project leader
Dr. Ernest Weingartner, co-leader
Dr. Bart Verheggen*, Julie Cozic, Günther Wehrle, Staffan Sjögren, Stefan van Ekeren, Dr. Martin Gysel,
Dr. M. Collaud Coen, MeteoSwiss, Payerne

*now at: Institute of Atmospheric and Climate Sciences, ETH Zürich

Project description:

Airborne aerosols affect our climate primarily by influencing the atmospheric energy budget through direct and indirect effects. Direct effects refer to the scattering and absorption of radiation and their influence on planetary albedo and the climate system. Indirect effects refer to the increase in available cloud condensation nuclei (CCN) due to an increase in anthropogenic aerosol concentration. This could lead to an increase in cloud droplet number concentration and a decrease in cloud droplet effective radius, when the cloud liquid water content (LWC) remains constant. The resulting cloud droplet spectrum could lead to reduced precipitation and increased cloud lifetime. The overall result would be an increase in cloud albedo which cools the Earth's climate. Despite the uncertainty, it is believed that in regions with high anthropogenic aerosol concentrations, aerosol forcing may be of the same magnitude, but opposite in sign to the combined effect of all greenhouse gases.

The Global Atmosphere Watch (GAW) program is an activity overseen by the World Meteorological Organization (WMO). It is the goal of GAW to ensure long-term measurements in order to detect trends and to develop an understanding of these trends. With respect to aerosols, the objective of GAW is to determine the spatio-temporal distribution of aerosol properties related to climate forcing and air quality up to multi-decadal time scales. Since the atmospheric residence time of aerosol particles is relatively short, a large number of measuring stations are needed. The GAW monitoring network consists of 23 Global (including now the Jungfrauoch, which was upgraded from a Regional to a Global station) and some 300 Regional stations. While Global stations are expected to measure as many of the key variables as possible, the Regional stations generally carry out a smaller set of observations.

The Jungfraujoch aerosol program is among the most complete ones worldwide. The current GAW instrumentation that is continuously run at the Jungfraujoch consists of

- CPC (TSI 3010) Particle number density (particle diameter $D_p > 10$ nm)
- Nephelometer (TSI 3563) Scattering coefficient at various wavelengths
- Aethalometer (AE-31) Absorption coefficient at various wavelengths; black carbon (BC) concentration
- MAAP Absorption coefficient; black carbon (BC) conc.
- Filter packs Aerosol major ionic composition (PM1 and TSP)
- Betameter and HiVol Aerosol mass (PM1 and TSP)

For these measurements, ambient air is sampled via a heated inlet (25 °C), designed to prevent ice build-up and to evaporate cloud particles at an early stage, ensuring that the cloud condensation nuclei and/or ice nuclei are also sampled. This is called the *total* inlet.

In warm months, the site is influenced by injection of planetary boundary layer air into the free troposphere during sunny afternoons due to thermal convection, while in winter it is usually in the undisturbed free troposphere. This causes the concentration of pollutants, including the aerosol loading, to be higher in summer than in winter (see Figure 1).




Figure 1. Long-term measurements of the light scattering coefficient (at 550 nm).

This data set was used to analyze the long-term trend of aerosol optical measurements at the Jungfraujoch (Collaud Coen *et al.*, submitted to J. Geophys. Res.). Since the aerosol variables are approximately lognormally distributed, the seasonal Kendall test and Sen's slope estimator were applied as non-parametric methods to detect the long-term trends for each month. The yearly trend was estimated by a least-mean square fit. The most significant trend is the increase (4-7% yr^{-1}) in light scattering coefficients during the September to December period. The light absorption and backscattering coefficients and the aerosol number concentration also show a positive trend during this time of the year. This increase of the extensive aerosol properties during autumn can tentatively be related either to the general temperature increase over Europe, which lengthens the period when the influence of the PBL is preponderant and increases the injection of PBL air masses into the lower FT over

large regions, or to long-range transport from areas with growing economy and increasing air pollution. The hemispheric backscattering fraction and the scattering exponent show an increase in size of small particles during most of the year, whereas the size of large particles remains constant. Generally the summer months at the Jungfraujoch, which are strongly influenced by planetary boundary layer air masses, do not show any long-term trend.

The fifth Cloud and Aerosol Characterization Experiment (CLACE 5) took place in February/March, 2006, with participation from different research groups. During this campaign, additional instrumentation was employed to characterize the aerosol size distribution (Scanning Mobility Particle Sizer, SMPS; Optical Particle Counter, OPC). The Max Plank Institute in Mainz (MPI) operated two Aerodyne Aerosol Mass Spectrometers (AMS) to measure the size segregated chemical composition. New in comparison with previous CLACE campaigns were the employment of an ice nucleus counter and an ATOF-MS for single particle analysis of ice nuclei by ETH Zürich.

Two additional inlets were used for these instruments: An *interstitial* inlet operated with a PM₂ cyclone impactor removed all cloud particles from the ambient air. Within a cloud the sampled air thus represents the interstitial (or unactivated) aerosol fraction. In addition, the Institute for Tropospheric Research (IfT) operated a Counterflow Virtual Impactor (CVI) (Mertes *et al.*, submitted to Environ. Sci. Technol.). The CVI was part of a new prototype sampling system (Ice-CVI) that allows for the separation of small ice particles from large ice crystals, cloud droplets and interstitial aerosol particles. The extracted ice particles are dried airborne in the system and the remaining residual particles which correspond to the former ice nuclei were analyzed with a variety of different instruments.

Differencing the response downstream of the different inlets provides insight in the fractionation of aerosol particles between the cloud phase and the interstitial phase. The activated fraction is defined as the fraction of the total particle number ($D_p > 100$ nm) that is activated into cloud droplets (obtained from total minus interstitial). Based on more than 900 hours of in-cloud measurements from winter and summer 2004 and winter 2005, this activated fraction has been related to several environmental factors such as liquid water content, number concentration of particles, temperature, and ice mass fraction of the cloud. These analyses revealed that the black carbon (BC) component of the ambient aerosol is activated into cloud droplets to the same extent as the bulk aerosol. Such a behaviour is not expected for freshly emitted soot particles because they have a hydrophobic nature. The soot particles on the Jungfraujoch experienced aging processes which transformed them into an internally mixed hygroscopic aerosol (Cozic *et al.*, 2006).

During wintertime this activated fraction is generally low (below 20%), because the presence of ice crystals causes liquid droplets to evaporate, thus transforming cloud droplets back into interstitial aerosol particles, as described by the Wegener-Bergeron-Findeisen process. When the cloud exists almost exclusively of liquid droplets (i.e. ice mass fraction approaching zero), the activated fraction increases to similar values as those encountered in summer (approximately 50%) (Verheggen *et al.*, submitted to *J. Geophys. Res.*). This is shown in Figure 2.




Figure 2. Activated fraction of particles as a function of ice mass fraction in the cloud. Circles denote the average, while horizontal stripes denote the 25 and 75 percentile values.

A similar analysis was performed for the partitioning of the black carbon (BC) mass concentration in mixed phase clouds. The BC mass concentration experiences a similar trend as the one observed for the bulk aerosol at low and intermediate ice mass fractions (Figure 2). A significant difference is found for the scavenged BC fraction at high ice mass fraction: The scavenged BC fraction is ~10% at ice mass fractions > 0.9 whereas the scavenged fraction of the bulk aerosol volume decreases to near-zero (see Figure 2). This indicates that the BC mass concentration is enriched in the ice phase compared to the bulk aerosol. This hypothesis is corroborated by the comparison of the BC mass fraction measured in the ice residuals with the BC mass fraction in the bulk aerosol (Figure 3). The fact that most points are located above the 1:1 line indicates that BC containing particles are enriched in the ice residuals (Cozic *et al.*, manuscript in preparation). This result is highly important for climate modeling since BC has a predominately anthropogenic origin.




Figure 3. Simultaneously measured BC mass fraction for ice residuals and for the bulk aerosol. The BC mass fraction is defined as the ratio of the particle BC mass concentration to total aerosol mass concentration.

Incorporation of the observed relation between number of ice crystals, particle number concentration and ice mass fraction into a global climate model suggests that the Wegener-Bergeron-Findeisen mechanism may have a dampening effect on the indirect effect of aerosols on climate (Weingartner *et al.* (2006), submitted).

Key words:

Atmospheric aerosol particles, aerosol-cloud interactions, aerosol climatic effects, radiative forcing, cloud condensation nuclei, hygroscopic growth, particle nucleation

Internet data bases:

<http://www.psi.ch/gaw>

<http://www.psi.ch/lac>

<http://aerosolforschung.web.psi.ch>

http://www.meteoschweiz.ch/web/en/climate/global_climate_monitoring/GAW_CH_Allg/GAW-aerosol.html

Collaborating partners/networks:

Dr. P. Viatte, MeteoSwiss, Payerne

Dr. C. Hüglin and Dr. S. Reimann, EMPA, Dübendorf

Prof. H. Burtscher, Institut für Sensoren und Signale der Fachhochschule Aargau (FHA), Windisch

Prof. U. Lohmann and Prof. T. Peter, Institute for Atmospheric and Climate Science, ETH Zürich

Prof. J. Heintzenberg, Institut für Troposphärenforschung, Leipzig, Germany

Dr. A. Petzold, Institute of Atmospheric Physics, DLR Oberpfaffenhofen, Germany

Prof. H. Coe and Prof. T. Choularton, School of Earth, Atmospheric and Environmental Sciences (SEAES), University of Manchester, Manchester, England

Dr. J. Schneider and Prof. S. Borrmann, University of Mainz, Particle Chemistry Department, Mainz, Germany

Prof. S. Weinbruch, Universität Darmstadt, Institut für Mineralogie, Darmstadt, Germany

Prof. M. Kulmala, Department of Physics, University of Helsinki, Helsinki, Finland

Dr. E. Fries, J. W. Goethe University, Institute for Atmosphere and Environment, Frankfurt, Germany

Scientific publications and public outreach 2006:

Refereed journal articles

Cozic, J., B. Verheggen, S. Mertes, P. Connolly, K. N. Bower, A. Petzold, U. Baltensperger, and E. Weingartner, Scavenging of black carbon in mixed phase clouds at the high alpine site Jungfraujoch, *Atmos. Chem. Phys. Discuss.*, 6, 11877–11912 (2006).

Conference papers

Baltensperger, U. and E. Weingartner, *Climate relevant aerosol research at the high altitude research station Jungfraujoch, Switzerland*, Proc. 75th Anniversary of the High Altitude Research Station Jungfraujoch, Interlaken, Switzerland, September 11-14 (2006).

Collaud Coen M., E. Weingartner, S. Nyeki, U. Baltensperger, *Long-term trend analysis of aerosol parameters at the Jungfraujoch*, Proc. 4-8 April 2006, EGU 2006, Vienna, Austria, April 4-8 (2006).

Collaud Coen M., E. Weingartner, U. Baltensperger, *Missing effects between the planetary boundary layer and the lower free troposphere detected by the measurement of aerosol parameters at the Jungfraujoch*, Proc. 8th International Congress on Aerobiology, Neuchâtel, Switzerland, August 21-25 (2006).

Collaud Coen M., E. Weingartner, S. Nyeki, U. Baltensperger, *Variability and long-term trends of aerosol parameters at the Jungfraujoch*, Proc. 75th Anniversary of the High Altitude Research Station Jungfraujoch, Interlaken, Switzerland, September 11-14 (2006).

Collaud Coen M., E. Weingartner, D. Schaub, C. Hueglin and U. Baltensperger, *Saharan dust events at the Jungfraujoch: a new detection method and a five-year climatology*, Proc. 75th Anniversary of the High Altitude Research Station Jungfraujoch, Interlaken, Switzerland, September 11-14 (2006).

Cozic, J., *The fate of black carbon in the atmosphere: Its incorporation into cloud droplets and ice crystals*, Proc. ASEFI 2006, Arcachon, France, October (2006).

Cozic, J., B. Verheggen, U. Baltensperger, E. Weingartner, *The fate of black carbon in the atmosphere: Rapid removal by wet deposition after aging*, Proc. 10th ETH-Conference on Combustion Generated Nanoparticles, Zürich, Switzerland, August 12-15, (2006).

Cozic, J., B. Verheggen, S. Mertes, P. Connolly, K. Bower, A. Petzold, U. Baltensperger, E. Weingartner, *Black carbon contribution to the aerosol phase and its scavenged fraction in mixed phase clouds at the high alpine site Jungfraujoch (3580m*

asl), Proc. 75th Anniversary of the High Altitude Research Station Jungfrauoch, Interlaken, Switzerland, September 11-14 (2006).

Cozic, J., B. Verheggen, S. Mertes, M. Flynn, P. Connolly, K. Bower, A. Petzold, U. Baltensperger, E. Weingartner, *Black carbon contribution to the aerosol phase and its scavenged fraction in mixed phase clouds at the high alpine site Jungfrauoch (3580m asl)*, Proc. 7th International Aerosol Conference, St. Paul, Minnesota, USA, 2, 1393, September 10-15 (2006).

Mertes S., B. Verheggen, S. Walter, M. Ebert, P. Connolly, J. Schneider, K.N. Bower, J. Cozic, A. Worringer, E. Weingartner, *Counterflow virtual impactor based collection of small ice particles in mixed-phase clouds for the physico-chemical characterisation of tropospheric ice nuclei*, Proc. 7th International Aerosol Conference, St. Paul, Minnesota, USA, 2, 1363, September 10-15 (2006).

Schneider J., S. Walter, J. Curtius, S. Borrmann, S. Mertes, E. Weingartner, B. Verheggen, J. Cozic, U. Baltensperger, *Chemical composition measurements of ice nuclei in mixed phase tropospheric clouds during the Cloud and Aerosol Characterization Experiments CLACE*, Proc. 7th International Aerosol Conference, St. Paul, Minnesota, USA, 2, 1356, September 10-15 (2006).

Vana M., A. Hirsikko, E. Tamm, P. Aalto, M. Kulmala, B. Verheggen, J. Cozic, E. Weingartner, U. Baltensperger, *Characteristics of air ions and aerosol particles at the high alpine research station Jungfrauoch*, Proc. 7th International Aerosol Conference, St. Paul, Minnesota, USA, 2, 1427, September 10-15 (2006).

Verheggen B., J. Cozic, E. Weingartner, M. Vana, P. Aalto, A. Hirsikko, M. Kulmala, U. Baltensperger, *Observations of atmospheric nucleation events in the lower free troposphere*, Proc. 7th International Aerosol Conference, St. Paul, Minnesota, USA, 2, 1415, September 10-15 (2006).

Verheggen B., E. Weingartner, J. Cozic, M. Vana, P. Aalto, A. Hirsikko, M. Kulmala, U. Baltensperger, *Nucleation events at a high alpine site: Particle growth and nucleation rates*, Proc. Formation and Growth of Atmospheric Aerosols, Minneapolis, September 8-9 (2006).

Weingartner E., *Erkenntnisse aus dem GAW-Aerosolprogramm am Jungfrauoch*, Proc. Feinstaub in der Schweiz, EMPA Dübendorf, January 20 (2006).

E. Weingartner, *Verhalten von Aerosolen in der realen Atmosphäre*, Proc. München, HDT Seminar, Minimierung der Partikelemissionen von Dieselfahrzeugen May 15-17 (2006).

Weingartner E., B. Verheggen, J. Cozic, M. Gysel, S. Sjogren, J. Duplissy, U. Baltensperger, U. Lohmann, S. Mertes, K.N. Bower, M. Flynn, P. Connolly, J. Crosier, J. Crosier, M. Gallagher, H. Coe, T. Choulaton, S. Walter, J. Schneider, J. Curtius, S. Borrmann, A. Petzold, M. Ebert, M. Innerle-Hof, A. Worringer, S. Weinbruch, E. Fries, E. Starokozhev, W. Püttmann, W. Jaeschke, M. Vana, A. Hirsikko, E. Tamm, P. Aalto, M. Kulmala, *Aerosol-cloud interactions in the lower free troposphere as measured at the high alpine research station Jungfrauoch in Switzerland*, Proc. 7th International Aerosol Conference, St. Paul, Minnesota, USA, 2, 1383, September 10-15 (2006).

Address:

Laboratory of Atmospheric Chemistry
Paul Scherrer Institut (PSI)
CH-5232 Villigen
Switzerland

Contacts:

Ernest Weingartner

Tel: +41 56 310 2405

Fax: +41 56 310 4525

e-mail: ernest.weingartner@psi.ch

Urs Baltensperger

Tel: +41 56 310 2408

Fax: +41 56 310 4525

e-mail: urs.baltensperger@psi.ch